

**MAGNITUDE AND EXTENT
OF SEDIMENT TOXICITY IN
THE HUDSON-RARITAN
ESTUARY**

Published @ 2017 Trieste Publishing Pty Ltd

ISBN 9780649118076

Magnitude and extent of sediment toxicity in the Hudson-Raritan Estuary by Various

Except for use in any review, the reproduction or utilisation of this work in whole or in part in any form by any electronic, mechanical or other means, now known or hereafter invented, including xerography, photocopying and recording, or in any information storage or retrieval system, is forbidden without the permission of the publisher, Trieste Publishing Pty Ltd, PO Box 1576 Collingwood, Victoria 3066 Australia.

All rights reserved.

Edited by Trieste Publishing Pty Ltd.
Cover @ 2017

This book is sold subject to the condition that it shall not, by way of trade or otherwise, be lent, re-sold, hired out, or otherwise circulated without the publisher's prior consent in any form or binding or cover other than that in which it is published and without a similar condition including this condition being imposed on the subsequent purchaser.

www.triestepublishing.com

VARIOUS

**MAGNITUDE AND EXTENT
OF SEDIMENT TOXICITY IN
THE HUDSON-RARITAN
ESTUARY**

Magnitude and Extent of Sediment Toxicity in the Hudson-Raritan Estuary

Edward R. Long

*National Ocean Service, Office of Ocean Resources Conservation
and Assessment, Seattle, WA*

Douglas A. Wolfe

*National Ocean Service, Office of Ocean Resources Conservation
and Assessment, Silver Spring, MD*

K. John Scott and Glen B. Thursby

*Science Applications International Corporation
Narragansett, RI*

Eric A. Stern

*U.S. Environmental Protection Agency
New York, New York*

Carol Peven

*Battelle Ocean Sciences
Duxbury, MA*

Ted Schwartz

*National Biological Survey
Washington, D.C.*

Table of Contents

List of Tables	i
List of Figures	iii
Abstract	1
I. Introduction	2
Contaminant Concentrations	4
Potential for Toxicant Effects	5
Previously Measured Biological Effects	8
II. Methods and Materials	15
Sampling Methods	19
Sediment Testing Methods	20
Estimates of the Spatial Extent of Toxicity	22
Chemical Analyses: Phase 1	23
Chemical Analyses: Phase 2	25
Data Analyses	27
III. Results	28
Solid-Phase Amphipod Tests	28
Elutriate/Liquid Phase Bivalve Larvae Tests	38
Microbial Bioluminescence Tests of Organic-Extracts	50
Polychaete and Sand Dollar Growth Tests	57
Estimates of Spatial Extent of Toxicity	58
Concentrations and Distribution of Contaminants in Sediments: Phase 1	60
Concentrations and distribution of Contaminants in Sediments: Phase 2	62
Relationships Between Toxicity and Physical-Chemical Parameters: Phase 1	76
Relationships Between Toxicity and Physical-Chemical Parameters: Phase 2	96
IV. Discussion	111
Incidence and Severity of Toxicity	111
Spatial Extent of Toxicity	117
Spatial Patterns in Toxicity	117
Correlations Among Toxicity Tests	121
Summary of Chemistry/Toxicity Relationships	122
V. Conclusions	126
Potential for Toxicity	126
Incidence of Toxicity	126
Spatial Patterns in Toxicity	127
Spatial Extent of Toxicity	127
Chemistry/Toxicity Relationships	127
Acknowledgments	128
References	129
Appendices	134

List of Tables

1.	Regions of the Hudson-Raritan Estuary in which the concentrations of selected toxicants in sediments exceeded respective effects ranges of Long and Morgan (1990). Adapted from Squibb et al. (1991)	6
2.	Spearman-rank correlations (Rho) between percent survival of <i>A. abdita</i> (n=9) and the concentrations of trace metals normalized to dry wt., aluminum and total organic carbon (TOC). From U.S. EPA EMAP monitoring data, 1990 (Schimmel et al., 1994)	11
3.	Mean percent mortality of <i>Eohaustorius</i> estuaries in sediments from Arthur Kill, Kill van Kull, and Newark Bay (from Aqua Survey, Inc., 1990a, 1990b)	12
4.	Locations of sites in the Hudson-Raritan Estuary sampled during Phase 1	15
5.	Locations of stations in Newark Bay and vicinity sampled during Phase 2	18
6.	Mean percent survival of <i>A. abdita</i> in 10-day solid-phase toxicity tests of sediments from the Central Long Island Sound (CLIS) control site (n = 5), 117 sampling stations (n = 5) and 39 sites (n = 3) and of <i>Diporeia</i> spp. in 9 samples from the Hudson-Raritan Estuary	31
7.	Mean percent amphipod (<i>A. abdita</i>) survival in the 1993 Newark Bay survey performed during Phase 2	36
8.	Mean percent survival and normal morphological development (expressed as percent of controls) in 48-hour tests of elutriates with the larvae of <i>Mulinia lateralis</i>	41
9.	Results of Microtox™ tests of microbial bioluminescence in organic extracts of sediments; mean EC50's (n = 2) and 95% confidence intervals for stations, and mean EC50's (n = 3) for sites	50
10.	Results of microbial bioluminescence (Microtox™) tests of sediments from the Hudson-Raritan estuary performed with three kinds of sediment extracts (from DeMuth et al., 1993)	57
11.	Results of polychaete (<i>Armandia brevis</i>) impaired growth tests, and sand dollar (<i>Dendraster excentricus</i>) impaired growth tests of sediment from the Hudson-Raritan estuary (from Rice et al., in press)	58
12.	Estimates of the spatial extent of toxicity* (km ² and percent of total area) in the Hudson-Raritan Estuary based upon the cumulative distribution functions of data from each of four test end-points	59
13.	Estimates of concordance in the spatial extent of toxicity* (km ² and percent of total area) in the Hudson-Raritan estuary among the four toxicity test end-points	59
14.	Estimates of the spatial extent of toxicity* (km ² and percent of total area) in Newark Bay and vicinity, based upon the cumulative distribution function of data from amphipod survival tests	60
15.	Concentrations of TCDD-equivalents (pg/g) in whole extracts and extract fractions determined in H4IIE rat hepatoma bioassays of sediments from Newark Bay	75
16.	Spearman-rank (rho, corrected for ties) correlations between dioxin equivalents determined in chemical analyses and dioxin equivalents (tcdd-eqs) determined in rat hepatoma bioassays of sediment extracts	76

17.	Spearman-rank correlations (Rho, corrected for ties) between four toxicity end-points (as percent of controls) and the concentrations of trace elements in Hudson-Raritan estuary sediments (n=38)	80
18.	Spearman-rank correlations (Rho, corrected for ties) between four toxicity end-points (as percent of controls) and the concentrations of acid-volatile sulfides (AVS) and simultaneously extracted trace metals (SEM) in Hudson-Raritan Estuary sediments (n=38)	80
19.	Spearman-rank correlations (Rho, corrected for ties) between four toxicity end-points (as percent of controls) and the concentrations of PCBs and pesticides in Hudson-Raritan Estuary sediments (n=38)	81
20.	Spearman-rank correlations (Rho, corrected for ties) between four toxicity end-points (as percent of controls) and the concentrations of PAHs in Hudson-Raritan Estuary sediments (n=38)	82
21.	Samples from the Hudson-Raritan estuary (Phase 1) stations that equalled or exceeded the respective ERM or SQC guideline concentrations for each major substance or class of compounds. Stations in which the concentration exceeded the guideline by >2x are listed in bold (n=38).....	83
22.	Average trace metal concentrations (ppm, dry wt. or $\mu\text{mole/g} \pm \text{s.d.}$) in samples that were not toxic, significantly toxic ($p < 0.05$), and highly toxic (percent survival <80% of controls) in amphipod tests, ratios between the averages, and ratios of highly toxic averages to SQGs	89
23.	Average pesticide and PCB concentrations in samples that were not toxic, significantly toxic, and highly toxic in the amphipod tests, ratios between the averages, and ratios of highly toxic averages to SQGs	90
24.	Average PAH concentrations in samples that were not toxic, significantly toxic, and highly toxic in the amphipod tests, ratios between the averages, and ratios between highly toxic averages and respective SQGs	92
25.	Average trace metal concentrations in samples that were not toxic, significantly toxic, and highly toxic in microtox tests, ratios between the averages, and ratios of highly toxic averages to SQGs	93
26.	Average pesticide and PCB concentrations in samples that were not toxic, significantly toxic, and highly toxic in the Microtox tests, ratios between the averages, and ratios of highly toxic averages to SQGs	94
27.	Average PAH concentrations in samples that were not toxic, significantly toxic, and highly toxic in the Microtox tests, ratios between the averages, and ratios between highly toxic averages and respective SQGs	95
28.	Spearman-rank correlations between percent amphipod survival and the concentrations of total trace metals and with the ratios of simultaneously extracted metals to acid-volatile sulfides in Phase 2 sediments	96
29.	Spearman-rank correlations between percent amphipod survival and the concentrations of chlorinated organic compounds in Newark Bay	97
30.	Spearman-rank correlations between percent amphipod survival and the concentrations of chlorinated dibenzo dioxin and dibenzo furan compounds in Newark Bay sediments	98
31.	Spearman-rank correlations between percent amphipod survival and the concentrations of polynuclear aromatic hydrocarbons in Newark Bay sediments	99

32.	Samples from the Phase 2 stations that equalled or exceeded the respective ERM or SQC values for each major substance or class of compounds	100
33.	Average concentrations of 2,3,7,8-tcdd and total cumulative dioxin TEQs in highly toxic and nontoxic samples from Newark Bay, ratios between the averages, and ratios between the highly toxic averages and the respective SQG	108
34.	Average concentrations of pesticides and PCBs in highly toxic and nontoxic samples from Newark Bay, ratios between the averages, and ratios between the highly toxic averages and the respective SQGs	108
35.	Average concentrations of total extractable and AVS simultaneously extracted trace metals in highly toxic and nontoxic samples from Phase 2, ratios between the averages, and ratios between the highly toxic averages and the respective SQGs	110
36.	Average concentrations of PAHs in highly toxic and nontoxic samples from Phase 2, ratios between the averages, and ratios between the highly toxic averages and the respective SQGs	111
37.	Summary of toxicity test results for each station and site sampled during Phase 1	112
38.	Summary of the numbers of Phase 1 stations and sites indicated as significantly toxic and numerically significant in each of four sediment toxicity test endpoints.....	116
39.	Spearman rank correlation coefficients for the four toxicity test end-points tested in Phase 1 as percent of controls	121
40.	Summary of toxicity/chemistry relationships for those chemicals that were significantly correlated with toxicity in the Phase 1 samples	123
41.	Summary of toxicity/chemistry relationships for those chemicals that were significantly correlated with toxicity in the Phase 2 samples	125