REPORT ON THE HERBACEOUS FLOWERING PLANTS OF MASSACHUSETTS, ARRANGED ACCORDING TO THE NATURAL ORDERS OF LINDLEY, AND ILLUSTRATED CHIEFLY BY POPULAR DESCRIPTIONS OF THEIR CHARACTER. PROPERTIES. AND USES

Published @ 2017 Trieste Publishing Pty Ltd

ISBN 9780649113156

Report on the herbaceous flowering plants of Massachusetts, arranged according to the natural orders of Lindley, and illustrated chiefly by popular descriptions of their character, properties, and uses by Chester Dewey

Except for use in any review, the reproduction or utilisation of this work in whole or in part in any form by any electronic, mechanical or other means, now known or hereafter invented, including xerography, photocopying and recording, or in any information storage or retrieval system, is forbidden without the permission of the publisher, Trieste Publishing Pty Ltd, PO Box 1576 Collingwood, Victoria 3066 Australia.

All rights reserved.

Edited by Trieste Publishing Pty Ltd. Cover @ 2017

This book is sold subject to the condition that it shall not, by way of trade or otherwise, be lent, re-sold, hired out, or otherwise circulated without the publisher's prior consent in any form or binding or cover other than that in which it is published and without a similar condition including this condition being imposed on the subsequent purchaser.

www.triestepublishing.com

CHESTER DEWEY

REPORT ON THE HERBACEOUS FLOWERING PLANTS OF MASSACHUSETTS, ARRANGED ACCORDING TO THE NATURAL ORDERS OF LINDLEY, AND ILLUSTRATED CHIEFLY BY POPULAR DESCRIPTIONS OF THEIR CHARACTER, PROPERTIES, AND USES

Trieste

M 3

REPORTS

ON THE

HERBACEOUS PLANTS

AND ON THE

QUADRUPEDS

OF

MASSACHUSETTS.

PUBLISHED AGREEABLY TO AN ORDER OF

THE LEGISLATURE,

BY THE COMMISSIONERS ON THE ZOOLOGICAL AND BOTANICAL SURVEY OF THE STATE,

CAMBRIDGE:

FOLSOM, WELLS, AND THURSTON, PRINTERS TO THE UNIVERSITY.

1640.

REPORT

277 0.6

ON THE

HERBACEOUS FLOWERING PLANTS

OF

MASSACHUSETTS,

ARRANGED ACCORDING TO THE NATURAL ORDERS OF LINDLEY,

AND

ILLUSTRATED CHIEFLY BY POPULAR DESCRIPTIONS

OF THEIR

CHARACTER, PROPERTIES, AND USES.

PUBLISHED AGREEABLY TO AN ORDER OF

THE LEGISLATURE,

BY THE COMMISSIONEES ON THE ZOOLOGICAL AND BOTANICAL SURVEY OF THE STATE,

CAMBRIDGE:

FOLSOM, WELLS, AND THURSTON, PRINTERS TO THE UNIVERSITY.

1840.

.

INTRODUCTORY REMARKS,

Page 5

CLASS I. VASCULARES. Flowering Plants.

82

12

r,

•

SUE-CLASS I. EXOGENÆ. Dicotyledones. TRIBE I. ANGIOSPERMÆ.

1. Polypetalous Plants.

Order	1.	Araliacen, .		23	23444	2010-023 222		25		12		13
Cider	2.	Umbellifera,	- 25	<i></i>	14		R.		1		2	15
	3.	Ranunculacea,	÷.	π	62		00					21
	4.	Papaveracea,	90	20	34	85	•	te	a.	1		29
	5.	Nymphreacere,		92		2		40		12		31
	7.	Hydropeltideæ,	32		22		2		32		73	32
	S.	Podophyllew,		20								32
	9.	Crucifera,	12		19		+2		-		65	33
	10.	Fumariacee,		43		94		- 83		38		39
	11.	Capperidea,	12		12		20		2			40
	22,	Berberidere,		20		53		•		2		41
	24.	Malvacea,	÷E		90		102		18		10	41
	36.	Hypericinese,		ie.		(ji		¥.)				44
	38.	Saxifragea,	(\mathbf{x})		12		47		12		\$3).	44
	46.	Caetere, .		20		3		20		$\overline{\mathcal{X}}$		46
	47.	Onagrariæ,	80		12		23		35		tS.	46
	48.	Haloragea,		-		3		80		×		48
	49.	Circæaceæ,	2 7		3		$\hat{\mathbf{u}}_{i}$		24		12	49
	51.	Loascæ, .				.*		8		÷.		49
	52.	Salicariæ,	30		32		22		2		22	50
	54.	Melastomaceæ,		32				0		54		51

-												Page
Order	62.		•		3		•0		Эć		2	51
	64.	Santalacete,		•		12.2		ē.		э÷		52
	72.		•		1		10		1			53
	73.	Rosaceæ, .		± 1		1		72		25		54
	77.	Leguminosæ,	0		13		1				35	60
	78.	Unicere, .		${\bf s}(t)$		1		${\bf k}({\bf i})$		$^{\circ}$		69
	88.	Euphorbiaceæ,			84		¥0		32		.	71
	116.	Rutaceæ, .		X						٠		72
	122.	Geraniaceæ,	20		35		32		83		* 3	73
	123.	Oxalideæ, .		32		88		\mathbf{x}		×		74
	124.	Tropxolex,	2		23		\$ 2		82		-22	75
	126.	Balsamineæ,				34 1		3		2		75
	129.	Polygaleæ,					85		25		* 3	76
	130.	Violaceæ, .				10		¥3				77
	131.	Passifloreæ,	86		1		$ \hat{X} $		24		37	80
	134.	Cistineæ, .				2.5		2				80
	136.	Sarraceniere,			st:		15		1			81
	137.	Droseracem,		26		0		30		33		82
	139.	Lineæ,	22		$\widetilde{\mathbf{C}}$		ii.		267		*	83
	140.	Caryophylleæ,		2		13						84
	143.	Elatineæ,			58		27				÷	91
	144.	Portulaceæ,		28		6		20				91
	147.	Crassulaceæ,	5		-						æ	93
	148.	Ficoidere, .		14				12		(Se)		94
	150.	Illecebrete,			÷		32		25		4	94
	151.	Amaranthaceæ,				35						95
	152.	Scleranthere,	63		18		£1€		•			96
	153.	Chenopodea,		1		42		33		÷.		96
	154,	Phytolaccew,	2		4				43		22	100
	156.	Polygoneæ,		1		:40						101
	158,	Nyctagineze,	¥.0		- 36		3.8		35			104
	163.	Podostemere,		2		æ		15		83		105
	164.	Callitrichineæ,	÷3		4		84		÷.		2	105
	165.	Ceratophyllem,		8						23		105
		2. Mo.	nop	etal	ous	Pla	nts.					
	173.	Pyrolaceæ, .		8		4		02		4		106
	174.	Campanulaceæ,			5 190							108
	175.	Lobeliaceæ,		1		æ		- 29				110

Order	181	Cucurbitacese,							æ			Pago 111
order	182.	Plantagineæ,			•	6		8	240	Ş.		114
	183.	Plumbagineæ,	x:	57		1		(•).	22	81		115
	184.	Dipsaceae,	51d	13	1	æ	186		225		:0	116
	186.	Composita,	2 5		12		245		78		94	116
	189.	Stellate, .		1		3		823		72		143
	195.	Asclepiader,	ŧ0				10		12			145
	196.	Apocyneae, .		23		-		nei		÷		146
	197.	Gentianea,	20		Si.		10		35		823	147
	198.	Spigeliacea,		ŝ				847) 1413		\mathcal{L}		149
	199.	Convolvulaceæ,			24		4.5		13		ses.	150
	200.	Polemoniaceæ,		83		÷		16		30		151
	207.	Primulaceæ,	83						35		S	152
	208.	Lentibulariæ,		1		4		321		\overline{M}		155
	210.	Orobancheæ,			25		te					156
	211.	Scrophularinea,		82		æ		÷		$(\mathbf{\hat{r}})$		157
	212.	Rhinanthaccæ,	÷.		33		\tilde{M}					163
	213.	Solaneæ, .				22		$\langle i \rangle$		(0)		164
	215.	Pedalineae,	×.		15		57		:12			172
	220.	Verbenaceæ,		83		\sim		$\overline{\mathcal{D}}$		30		172
	221.	Labiatæ,	12		33		\hat{x}		÷		÷.	174
	222.	Boraginere,		20		\sim		121		¥.		184
	226.	Hydrophylleæ,					12		37			187
		TRIBE II.	G3	r MD	os	ų,		187				

SUB-CLASS II. ENDOGENÆ. Monocotyledones.

-

TRIBE I. PETALOIDE,E.

Order 229.	Alismaceæ,						20		æ		188
231.	Hydrocharideæ,					83		28		÷	190
232.	Commelinew,		85		$(\mathbf{k}$		¥0				191
233.	Xyrideæ,	ξį.		8		\mathbf{v}		S_{4}^{*}			191
235.	Hypoxideæ,		85		۰		25				192
238.	Amaryllidex,	×.		17				10		12	193
239.	Irideæ, .		83		12		£ΰ		$\overline{(2)}$		194
240.	Orchideæ,	90		8		ý.		S.;		i.	196
244.	Junceæ, .		2				ŧ		3		202
245.	Melanthaceæ,	8		S2		50		8		•	205

vii

Order 246.	Pontedereæ,			(43) (43)		ų,		12		23		Page 206
247.	Asphodelez,		es.		*		25		50			207
249.	Smilacew,					×		19		•2		210
250.	Dioscores,		(4)		36		4		20		18	214
251.	Liliacea,	÷.		3		1		82		1		215
253.	Restincoæ,				8		Q				S.	217
255.	Typhacea,			æ		30		38		25		218
256.	Aroidex,								8			219
258.	Fluviales,	÷		23		32		34		10		223
259.	Juncagine®,				4		3		83		2	225
260.	Pistiacow,			1.00				: •		20		226
	TRI	BE	п.	GI	UM	AC	EÆ.					
Order 261.	Gramineæ,					16				-		231
262.	Cyperacea,		ĸ		3		•		\times		٠	254
				-								
INDEX,	*	3		e		×		3				269

viii

REPORT

OS THE

HERBACEOUS PLANTS

0F

MASSACHUSETTS.

BY THE REV. CHESTER DEWEY,

PROCESSOR OF CHEMISTRY, BOTANY, AND NATURAL PHILOSOPHY IN THE BERKSHIRE NEDICAL INSTITUTION AT PITTSFIELD.