

**CATALOGUE OF THE SPECIMENS
OF HEMIPTERA HETEROPTERA
IN THE COLLECTION OF THE
BRITISH MUSEUM, PART VIII**

Published @ 2017 Trieste Publishing Pty Ltd

ISBN 9780649115440

Catalogue of the specimens of Hemiptera heteroptera in the collection of the British museum,
Part VIII by Francis Walker

Except for use in any review, the reproduction or utilisation of this work in whole or in part in
any form by any electronic, mechanical or other means, now known or hereafter invented,
including xerography, photocopying and recording, or in any information storage or retrieval
system, is forbidden without the permission of the publisher, Trieste Publishing Pty Ltd, PO Box
1576 Collingwood, Victoria 3066 Australia.

All rights reserved.

Edited by Trieste Publishing Pty Ltd.
Cover @ 2017

This book is sold subject to the condition that it shall not, by way of trade or otherwise, be lent,
re-sold, hired out, or otherwise circulated without the publisher's prior consent in any form or
binding or cover other than that in which it is published and without a similar condition
including this condition being imposed on the subsequent purchaser.

www.triestepublishing.com

FRANCIS WALKER

**CATALOGUE OF THE SPECIMENS
OF HEMIPTERA HETEROPTERA
IN THE COLLECTION OF THE
BRITISH MUSEUM, PART VIII**

22
523
Hem.
pt.
27

Library of
E. P. VAN DUZER

CATALOGUE

OF THE

SPECIMENS

OF

HEMIPTERA HETEROPTERA

IN

THE COLLECTION

OF THE

BRITISH MUSEUM.

PART VIII.

BY

FRANCIS WALKER.

PRINTED FOR THE TRUSTEES OF THE BRITISH MUSEUM:
LONDON, 1873.

LONDON I

E. NEWMAN, PRINTER, 9, DEVONSHIRE STREET, BISHOPSGATE.



P R E F A C E.

THE object of the present Catalogue is to give a complete List of all the genera and species of Hemiptera Heteroptera known to exist in the collections of European and American Entomologists. The letters *a*, *b*, *c*, &c., after the species, denote the specimens now contained in the British Museum, followed by the *habitat* and the mode in which each of them was obtained; and the absence of these *letters* indicates the species which are desiderata, and therefore desirable to be procured for the collection.

J. E. GRAY.

BRITISH MUSEUM,

August 1st, 1873.

CATALOGUE
OF
HEMIPTERA HETEROPTERA.

PART VIII.

Genus 6. OPINUS.

Tapeinus, *De Lap. Hem.* 81—*Opinus*, *De Lap. Hem. Suppl. Serv. Hist. Hem.* 338—*Tapeinus*, *Burm. Handb. Ent.* ii. 236.

America.

Div. 1.

Thymbreus, *Stal, Ofv. K. V. Ak. Forh.* xvii. 185.

The two following species were inserted in Pt. VII., p. 132, as *Pachynomi*.

1. *OPINUS PYRRHOPTERUS*.

Opinus pyrrhopterus, *Stal, Rio Jan. Hem.* 72—*Thymbreus pyrrhopterus*, *Stal, Ofv. K. Vet. Ak. Forh.* xvii. 186.

Brazil.

2. *OPINUS CROCINOPTERUS*.

Thymbreus crocinopterus, *Stal, Stett. Ent. Zeit.* xxiii. 457.
Mexico.

3. *OPINUS PYGOLEUS*.

Fulvus, fere linearis; caput prothoracis lobo antico paullo longius; antennarum articulus 2ns piecus, 3us 4usque testacei; prothoracis lobus anticus lobo postico ria brevior, anteice bituberculatus; femora anteice robusta, subtus spinulosa; ala anteice nigra basi fulva.

Tawny, nearly linear. Head small, short, a little longer than the fore lobe of the prothorax; ante-ocular part a little shorter than the post-ocular. Antennæ longer than the head and the prothorax together; first joint as long as the ante-ocular bead; second piecous, stout, twice as long as the first; third and fourth testaceous, very slender; third much shorter than the second; fourth as long as the third. Prothorax with a slight longitudinal furrow. Fore lobe nearly as long as the hind lobe, with a

small tubercle on each side of the fore border. Hind lobe with the hind angles rounded, not prominent. Scutellum blackish, equilaterally triangular. Fore femora stout, spinulose beneath. Fore wings black, tawny towards the base. Length of the body $2\frac{1}{2}$ lines.

a. Tejua. Presented by the Rev. H. Clark.

Africa.

4. OPINUS ACUTICEPS.

Niger, longissime oralis; caput prothoracis lobo antico triplo longius; prothorax rufus, lobis agnolongis; pedes rufi, tibis quatuor posterioribus femoribusque nigris, tibiis anticis nigro unifasciatis.

Black, dull, very elongate-oval. Head lanceolate, thrice as long as the fore lobe of the prothorax; ante-ocular part twice as long as the post-ocular. First joint of the antennæ stout, shorter than the head, much longer than the second. Prothorax red, with rounded and not prominent angles. Fore lobe as long as the hind lobe, convex on each side of the longitudinal furrow. Legs red; femora and four posterior tibiae black above; fore tibia with a black band. Length of the body $4\frac{1}{2}$ lines.

a. Sierra Leone. From Mr. Foxcroft's collection.

South Asia and Eastern Isles.

Div. 1.

5. OPINUS RUBROPICTUS.

tubropictus, Serv. Hist. Hem. 339.

Java.

Div. 2.

Sminthus, Stal, Ofv. K. Vet. Ak. Forh. xvii. 186.

6. OPINUS PROXIMUS.

proximus, Serv. Hist. Hem. 340—Sminthus proximus, Stal.

Java.

7. OPINUS PICTUS.

pictus, De Lap. Hém. 78, 82. Serv. Hist. Hem. 340—Tapinus pictus, Burm. Handb. Ent. ii. 236. H.-Sch. Wanx. Ins. viii. 56, pl. 267, f. 825—Hammatocerus pictus, Blanch. Hist. Nat. Ins. iii. 105—Sminthus pictus, Stal.

Java.

Var.? *proximus, Serv. Hist. Hem. 340.*

8. OPINUS INCONSPICUUS.

Tapirus inconspicuus, H.-Sch. Wanx. Ins. viii. 57, pl. 26, f. 826—Sminthus inconspicuus, Stal.

Java.

9. *OPINUS RUFUS.*

rufus, *De Lap. Hém.* 78, 82. *Serv. Hist. Hem.* 340—*Tapirus rufus*,
Burm. Handb. Ent. ii. 236.

a. Java. Presented by J. C. Bowring, Esq.
b, c. —?

10. *OPINUS GRATIOSUS.*

Sminthus gratiosus, *Stal, A. S. E. F. Acad. Sér.* iii. 55.
Tondano.

11. *OPINUS LIMBATICOLLIS.*

Sminthus limbaticollis, *Stal, Ofv. K. Vet. Ak. Forh.* xxiii. 238.
Malacca.

Div. 3.

Ienusus, *Stal, Ofv. K. Vet. Ak. Forh.* xvii. 186.

12. *OPINES PYRRHUS.*

Ienusus Pyrrhus, *Stal, Ofv. K. Vet. Ak. Forh.* xvii. 187.
Ceylon.

13. *OPINUS PUNCTORIUS.*

punctarius, *Stal, A. S. E. F. Acad. Sér.* iii. 55.
Wagion.

14. *OPINUS STRIGIFER.*

Fam. Rufus, *depressus*, *subfusiformis*, *subitus planus testaceus*; *caput subellipticum*, *prothoracis lobo antico equeolongum*; *antennae nigrae*; *prothoracis lobi aquilongi*; *abdomen apice nigricans*; *femora antica valida*; *membrana nigricans*.

Female. Red, flat, smooth, shining, fusiform, testaceous beneath. Head subelliptical, as long as the fore lobe of the prothorax; ante-ocular part as long as the post-ocular. Antennæ black, longer than the head and the prothorax together; first joint very short; second about six times as long as the first; third longer than the second. Prothorax with a longitudinal furrow; fore lobe as long as the hind lobe. Abdomen blackish at the tip; its disk beneath and the pectus flat. Fore femora stout. Membrane blackish. Length of the body $4\frac{1}{2}$ lines.

a. Malacca. Presented by W. W. Saunders, Esq.

15. *OPINUS RUGICOLLIS.*

Mas et fem. *Rufus*, *longissime ovalis*; *caput prothoracis lobo antico longius*; *antennæ nigrae*, *corporis dimidio longiores*; *prothoracis lobus anticus rugulosus*, *lobo postico paullo brevior*; *femora antica crassa*; *sulci tibiarum anticarum brevissimi*; *membrana nigra*.