WISCONSIN GEOLOGICAL AND NATURAL HISTORY SURVEY. BULLETIN NO. XII., SCIENTIFIC SERIES NO.3: THE PLANKTON OF LAKE WINNEBAGO AND GREEN LAKE

Published @ 2017 Trieste Publishing Pty Ltd

ISBN 9780649513710

Wisconsin Geological and Natural History Survey. Bulletin No. XII., Scientific Series No.3: The Plankton of Lake Winnebago and Green Lake by C. Dwight March

Except for use in any review, the reproduction or utilisation of this work in whole or in part in any form by any electronic, mechanical or other means, now known or hereafter invented, including xerography, photocopying and recording, or in any information storage or retrieval system, is forbidden without the permission of the publisher, Trieste Publishing Pty Ltd, PO Box 1576 Collingwood, Victoria 3066 Australia.

All rights reserved.

Edited by Trieste Publishing Pty Ltd. Cover @ 2017

This book is sold subject to the condition that it shall not, by way of trade or otherwise, be lent, re-sold, hired out, or otherwise circulated without the publisher's prior consent in any form or binding or cover other than that in which it is published and without a similar condition including this condition being imposed on the subsequent purchaser.

www.triestepublishing.com

C. DWIGHT MARCH

WISCONSIN GEOLOGICAL AND NATURAL HISTORY SURVEY. BULLETIN NO. XII., SCIENTIFIC SERIES NO.3: THE PLANKTON OF LAKE WINNEBAGO AND GREEN LAKE

Trieste

WISCONSIN GEOLOGICAL AND NATURAL HISTORY SURVEY.

E. A. BIRGE, Ph.D. Sc.D., Director.

BULLETIN NO. XII.

E E

-

K to c

ı,

SCIENTIFIC SERIES NO. 3.

THE PLANKTON

OF

Lake Winnebago and Green Lake

B¥

C. DWIGHT MARSH Professor of Biology in Ripon College.

> MADISON, WIS. PUBLISHED BY THE STATE. October, 1903.

Misconsin Geological and Ratural Distory Survey.

.

1

BOARD	OP	COMMISSIONERS.
-------	----	----------------

ROBERT M. LAFOLLETTE, Governor of the State. CHARLES B. VAN HISE, President, President of the University of Wisconsin. CHARLES P. CART, Vice-President, State Superintendent of Public Instruction. CALTERT SPENSLET. President of the Commissioners of Fisheries. JOEN J. DAVIS, Secretary, President of the Wisconsin Academy of Sciences, Arts, and Letters. STAFF OF THE SURVEY, E. A. BIRGE, Director of the Survey. 8. WEIDMAN, Geologist. Burvey of Central and Northern Wisconsin. U. S. GRANT, Goologist. Servey of Southwestern Wisconsin. N. M. FINNEMAN, Geologist. Physical Geography of Lake Region. C. D. MARSH, Biologist, Biology of Lakes. L. S. SETTE, Civil Engineer, Survey of Lakes and Elvers. W. D. SWITH, Field Assistant. E. T. HANCOCK, Field Assistant.

Consulting Geologist.

T. C. CHARBARLIN, Pleistocene Geology.

pi m

+ : :

.

CONTENTS.

	Page
CHAPTER I. OUTLINE OF STUDY	1-10
Lake Winnebago and Green Lake.	1
Methods of making collections	8
CHAPTER II. ANNUAL DISTRIBUTION OF THE ORGANISMS OF THE	
PLANETON	11-46
Distribution of C. brevispinosus and C. pulchellus in Wiscon-	
sin lakee	27
The "bloom"	36
Annual distribution of the total plankton	38
Constituents which produce plankton maxima	39
Comparison of amount of plankton in different years	41
Comparison of plankton in Green lake and Lake Winnebago	
Comparison with the plankton of other lakes	43
CRAPTER III. DISCUSSION OF RESULTS	47-59
Value of plankton collections	47
Relative importance of plankton constituents in producing	
maxima	49
Comparison of plankton and temperature curves	49
Comparison of curves of total plankton with curves of plankton	
contituents	51
Horizontal distribution	51
Comparison of plankton collections over mucidy and stony bot-	
toms	59
CHAPTER IV. DISTRIBUTION OF SPECIES	60-69
Geographical distribution	60
Comparison of the faunae and florae of different classes of	
lakes	62
Comparison of plankton of successive years	65
Relative value of deep and shallow lakes for the production of	
fish	67
LIST OF PARES OCCUP	70-71

162021

CONTENTS.

	Page
APPENDIX. STATISTICAL TABLES	73
Table I. Total volumes of plankton per square meter in Green	
lake	73
Table II. Total volumes of plankton per square meter in Lake	
Winnebago	74
Table III. Total volumes of plankton per square meter in va-	
rious Wisconsin lakes	75
Table IV. Numbers of D. minutus and D. sicilis per square	
meter in Green lake	77
Table V. Numbers' of D. oregonensis per square meter in	
Lake Winnebago	78
Table VI. Numbers of E. lacustris per square meter in Lake	
Winnebago	78
Table VII. Numbers of E. Jacustris per square motor in Green	
lake	79
Table VIII. Numbers of L. macrurus per square meter in	
Green lake	79
Table 1X. Numbers of C. brevispinosus per square meter in	
Lake Winnebago	80
Table X. Numbers of C. pulchellus per square meter in Lake	5223
Winnebego	80
Table XI. Numbers of C. teuckarti per square meter in Lake	-
Winnebago	81
Table XIL Numbers of C. prasinus per square meter in	
Green lako	81
Table XIII. Numbers of copepod larvae per square meter in	
Lake Winnebage Table XIV. Numbers of copeped larvae per square meter in	82
	00
Green lake Table XV. Nos. of <i>Diaphanosoma brachywrum</i> per square	82
에 가슴 바람에 있는 것을 만들었다. 것은 것은 것을 알았는 것을 하는 것을 것을 것을 것을 것을 것을 것을 하는 것을 하는 것을	00
meter in Lake Winnebago Table XVI. Numbers of D. hyatina per square meter in Lake	83
Winnebago	83
Table XVII. Numbers of D, hyaling per square meter in	00
Green lake	84
Table XVIII. Numbers of Bosmina per square meter in	01
Green lako	84
Table XIX, Numbers of Eurycercus lamellatus per square	
meter in Lake Winnebago	85
Table XX. Numbers of Chydorus per square meter in Lake	
Winnebago	85
Table XXI. Numbers of Leptodora per square meter in Lake	
Winnebago	86
Table XXII. Numbers of Cypris per square meter in Lake	50.52
Winnebago	86
Table XXIII, Maximum depth of Wisconsin lakes	87
INDEX	91
	1000

į.

Ľ

۱

iv

ILLUSTRATIONS.

53

4

PLATE	AGE
I. Curves of annual distribution of <i>Gloiotrichia</i> per square meter in Lake Winnebago	6
II. Curves of annual distribution of Diaptomus sicilis and	
Diaptomus minutus per square meter in Green lake . III. Curves of annual distribution of Diaptomus oregonensis	29
persquare meter in Lake Winnebago	23
IV. Curves of annual distribution of Epischura lacustris per square meter in Lake Winnebago	25
V. Curves of annual distribution of Epischura lacustris per	
square meter in Green lake	25
rus per square meter lo Green lake	26
VII. Curves of annual distribution of Cyclops brevispinosus per square meter in Lake Winnebago	26
VIII. Curves of [annual distribution of Cyclops breviepinosus	93802
per square meter in Green lake	26
square meter in Lake Winnebago	27
X. Curves of annual distribution of Cyclops Leuckarti per square meter in Green lake and Lake Winnebago	28
XI. Curves of annual distribution of Cyclops prasinus per	20
equare meter in Green lake	29
XII. Curves of annual distribution of copepod larvae per square meter in Green lake	29
XIII. Curves of annual distribution of copepod larvae per square	
meter in Lake Winnebago	29
chyurum per square meter in Lake Winnebago ,	31
XV. Curves of annual distribution of Daphnia hyalina per	01
square meter in Green lake	31
D. retrocurva per square meter in Lake Winebago .	79

ILLUSTRATIONS.

PLATE		PAGE
XVII.	Curves of annual distribution of Bosmina per square me-	
	ter in Green lake	33
XVIIL	Curves of annual distribution of Eurycercus lamellatus	
	per square meter in Lake Winnebago	34
XIX.	Curves of annual distribution of Chydorus sphaericus	
	per square meter in Lake Winnebago	35
XX.	Curves of annual distribution of Leptodora hyalina per	
	equare meter in Lake Winnebago	36
XXI.	Curves of annual distribution of total plankton in Lake	
	Winnebago per square meter	38
XXII.	Curves of annual distribution of total plankton in Green	
	lake per square meter	38

62

vī

THE PLANKTON OF LAKE WINNEBAGO AND GREEN LAKE.

CHAPTER I.

OUTLINE OF STUDY.

The attempt in this investigation has been to make a comparative study of the plankton of two lakes of different types, with a collateral study of such other lakes as could be easily reached.

It was thought that if, after a somewhat careful preliminary study, the plankton of the two lakes could be kept under observation for a considerable period of time, many facts in regard to the annual and goographical distribution of the animals and plants could be secured, and perhaps some generalizations could be reached in regard to the principles controlling such distribution.

LAKE WINNEBAGO AND GREEN LAKE.

Lake Winnebago was chosen for the type of the shallow lake mainly because of its importance from the standpoint of the production of fish. It is known to be an enormously productive body of water, and the fishing industry has employed a great many men in Oshkosh. Netting has been forbidden by legislative enactment for several years, and, in spite of many infraotions of the law, it is generally acknowledged among all but interested parties, that the number of fish has greatly increased in