ATLAS AND PRINCIPLES OF BACTERIOLOGY AND TEXT-BOOK OF SPECIAL BACTERIOLOGIC DIAGNOSIS; PART I - ATLAS

Published @ 2017 Trieste Publishing Pty Ltd

ISBN 9780649066872

Atlas and Principles of Bacteriology and Text-Book of Special Bacteriologic Diagnosis; Part I - Atlas by Dr. K. B. Lehmann & Dr. R. O. Neumann

Except for use in any review, the reproduction or utilisation of this work in whole or in part in any form by any electronic, mechanical or other means, now known or hereafter invented, including xerography, photocopying and recording, or in any information storage or retrieval system, is forbidden without the permission of the publisher, Trieste Publishing Pty Ltd, PO Box 1576 Collingwood, Victoria 3066 Australia.

All rights reserved.

Edited by Trieste Publishing Pty Ltd. Cover @ 2017

This book is sold subject to the condition that it shall not, by way of trade or otherwise, be lent, re-sold, hired out, or otherwise circulated without the publisher's prior consent in any form or binding or cover other than that in which it is published and without a similar condition including this condition being imposed on the subsequent purchaser.

www.triestepublishing.com

DR. K. B. LEHMANN & DR. R. O. NEUMANN

ATLAS AND PRINCIPLES OF BACTERIOLOGY AND TEXT-BOOK OF SPECIAL BACTERIOLOGIC DIAGNOSIS; PART I - ATLAS

Trieste

ATLAS AND PRINCIPLES

OF

BACTERIOLOGY

AND TEXT-BOOK OF

SPECIAL BACTERIOLOGIC DIAGNOSIS

BY

PROF. DR. K. B. LEHMANN Director of the Hygicalic Institute in Würzburg

AND

R. O. NEUMANN, DR. PHIL. and MED. Assistant in the Hygienic Institute in Würzburg

AUTHORIZED TRANSLATION FROM THE SECOND ENLARGED AND REVISED GERMAN EDITION

EDITED BY

GEORGE H. WEAVER, M.D. Assistant Professor of Pathology, Rush Medical College, Chicago

PART I-ATLAS

With 632 Figures on 69 Lithographic Plates

PHILADELPHIA AND LONDON W. B. SAUNDERS & COMPANY 1901

INDEX OF PLATES.

PLATE.

٠

- 1. Streptococcus pyogenes. Rosenbach.
- 2. Streptococcus lanceolatus. Gamaleia. (Diplococcus pneumoniæ. A. Fränkel.)

- Sarcina flava. De Bary, emended by Lehm, and Stubenrath.
 Sarcina aurantiaca. Flügge.
 Sarcina cervina. Stubenrath. Sarcina pulmonum. Virchow. Sarcina erythromyxa. Král. Sarcina lutea. Flügge. Sarcina aurantiaca. Flügge. Sarcina rosea. Schröter, emended by Zimmermann. Micrococcus badius. Lehm. and Neum.
- Sarcina canescens. Stubenrath. 6. Micrococcus luteus. Cohn, emended by Lehm. and Neum. Sarcina pulmonum. Virchow, Hauser.
- Micrococcus progenes a aureus. (Ros.) Lehm. and Neum. (Staphylococcus pyogenes aureus. Rosenbach.)
 Micrococcus pyogenes y albus. (Ros.) L. and N. (Staphylo-
- coccus pyogenes albus. Rosenbach). Micrococcus pyogenes β citreus. (Ros.) (Staphylococcus pyogenes citreus. Rosenbach.) Micrococcus candicans, Flügge. 10. Micrococcus gonorrhϾ. Neisser, Bumm. 11. Micrococcus roseus. (Bumm.) Lehm. and Neum. 12. Bacterium septicæmiæ hæmorrhagicæ. Hüppe. (Chicken chol-

- era, rabbit septicemia.)

- era, rabbit septicemia.)
 13. Bacterium pestis. Lehm. and Neum.
 14. Bacterium acidi lactici. "Hüppe. (Lactic acid bacillus.)
 15. Bacterium pneumoniæ. Friedländer.
 16. Bacterium typhi. Eberth, Gaffky. (Typhoid bacillus.)
 17. Bacterium typhi. Eberth, Gaffky.
 18. Bacterium coli. (Escherich.) Lehm. and Neum.
 19. Bacterium latericium. Adametz. Bacterium latericium. Adametz.
 19. Bacterium hæmorrhagicum. (Kolb.) Lehm. and Neum. (Morb. Werlhofii.)
- 21. Bacterium prodigiosum. (Ehrenberg.) Lehm. and Neum.

5

PLATE.

- 22. Bacterium kiliense. (Breunig and Fischer.) Lehm. and Neum.
- 23. Bacterium violaceum. (J. Schröter.) Lehm, and Neum.
- 24. Bacterium pyocyaneum. (Flügge.) Lehm. and Neum. (Green pus.)
- Bacterium fluorescens. (Flügge.) Lehm. and Neum. (Bacil-lus fluorescens liquefaciens, Flügge.)
- 26. Bacterium putidum. (Flügge.) Lehm. and Neum.
- 27. Bacterium syncyaneum. (Ehrenberg.) Lehm. and Neum. (Bacillus cyanogenes Flügge. Blue milk.) 28. Bacterium syncyaneum. (Ehrenberg.) Lehm. and Neum.
- 29. Bacterium Zopfii. Kurth.
- 30. Bacterium Zopfii. Kurth.
- 31. Bacterium vulgare. (Hauser.) Lehm. and Neum. (Proteus vulgaris Hauser.)
- 32. Bacterium vulgare β mirabilis. (Hauser.) Lehm, and Neum,
- 33. Bacterium erysipelatos suum. (Löffler.) Migula. (Hog erysipelas.)
 - Bacterium murisepticum. (Flügge.) Migula. (Mouse septicemia.)

- Bacillus anthracis.
 Bacillus anthracis.
 Bacillus anthracis.
 Bacillus anthracia.
 Cohn and R. Koch. (Splenic fever.)
 Bacillus anthracia.
 Cohn and R. Koch. (Splenic fever.)
- 37. Bacillus mycoides. Flügge. (Root-bacillus.)
- 38. Bacillus mycoides. Flügge.
- Bacillus butyricus. Huppe. (Butyric acid bacillus.) (Flögge.) Migula.
- Bacillus vulgatus.
- 39. Bacillus subtilis. F. Cohn. (Hay-bacillus.)
- 40. Bacillus subtilis. F. Cohn.
- 40. Bacillus subuns. F. Com. 41. Bacillus megatherium. De Bary. 41. Bacillus megatherium. (Flügge.) Migula. (B. mesentericus vul-Bacillus vulgatus. (Flügge.) Mia gatus Flügge.) Potato bacillus.
- 43. Bacillus mesentericus. (Flügge.) Lehm. and Neum. (B. mesentericus fuscus Flügge.)
- 44. Bacillus tetani. Nicolaier. (Tetanus bacillus, lock-jaw.)
- 45. Bacillus Chauvosi. Macè. (Symptomatic anthrax, black-leg.)
- Bacillus œdematis maligni.
 Vibrio choleræ. (Koch.) Koch.
- Buchner. (Comma bacillus,)
- 48. Vibrio choleræ. Buchner. (Koch.)
- 49. Vibrio choleræ. (Koch.) Buchner.
- 50. Vibrio choleræ. (Koch.) Buchner.
- 51. Vibrio cholerse. (Koch.) Buchner.
- Vibrio Metschnikovii. Gamaleia.
- 52. Vibrio Proteus. Buchner. (Vibrio Finkler.)
- Vibrio danubicus. Heider.
 - Vibrio berolinensis. Rubner. Vibrio aquatilis. Günther.
- 54. Vibrio albensis, Lehm, and Neum, (Phosphorescent Elbe Vibrio.)
- 55. Spirillum rubrum. v. Esmarch.

PLATE.

٠

- Spirillum concentricum. Kitasato. 56. Spirillum serpens. (E. O. Müller.) Zettnow. Spirilla from nasal mucus. Spirillum undula. Ehrenberg. Spirillum spermatozoides. Löffler, Spirochæte of oral mucus. Spirillum Obermeieri. F. Cohn. (Spirilla of recurrent fever.) 57. Corynebacterium malleï. (Löffler.) Lehm. and Neum. (Glanders bacillus.) 58. Corynebacterium diphtheriæ. (Löffler.) Lehm. and Neum. (Diphtheria bacillus.) Corynebacterium pseudodiphtheriticum. (Hoffmann-Wellenhof.) Lehm, and Neum. (Pseudodiphtheria bacillus.) Corynebacterium xerosis. (Kuschbert, Neisser.) Lehm. and Neum. (Xerosis bacillus.) Corynebacterium diphtheriæ, L. and N. Corynebacterium pseudodiphtheriticum, L. and N. Corynebacterium xerosis. L. and N. 60. Corynehacterium diphtheriæ. L. and N. Corynebacterium pseudodiphtheriticum. L. and N. Corynebacterium xerosis. L. and N. 61. Mycobacterium tuberculosis. (Koch.) L. and N. (Tubercle bacillus.) 62. Mycobacterium lepræ. (Arm. Hansen.) L. and N. (Lepra bacillus.) Mycobacterium tuberculosis y piscicola. L. and N. 63. Mycobacterium lacticola β perrugosum. L. and N. Mycobacterium phleï. Lehm. and Neum. Mycobacterium lacticola 4 planum. L. and N.
 Actinomyces bovis. Harz. (Actinomycosis.)
 Actinomyces farcinicus. (Nocard.) Gasperini. (Fracin du
- bœuf.)
- 67. Actinomyces chromogenes. Gasperini. (Cladothrix dichotoma Autorum non Cohn.)
- 68. Bacterium tussis convulsivæ. (Czaplewski and Hensel.) (Whooping-cough.)

Bacterium ulceris cancrosi. (Ducrey, Kruse.) L. and N. (Ulcus molle.)

Streptococcus meningitidis cerebrospinalis (Weichselbaum). Lehm. and Neum.


Bacterium influenze. (R. Pfeiffer.) L. and N. (Influenza bacillus.)

Bacillus gangrænæ pulpæ. Arkövy.

69. Leptothrix epidermidis. Biz.

Tab. 1.

-



ith. Anst F Reichhold, München,

- 12. si



Streptococcus pyogenes. Rosenbach.

I. Agar streak culture, ten days at 37°.

II. Gelatin stab culture, six days at 22°. So vigorous a growth does not often occur.

III. Agar stab culture, six days at 37°. Stab canal.

IV. Agar stab culture, six days at 37°. Surface growth.

V. Gelatin plate, six days at 22°.

VI. Gelatin plate, six days at 22° . \times 70. Somewhat abnormal form with irregular borders. The larger colonies are superficial; the smaller, deep.

VII. Gelatin plate, six days at 22° . \times 70. Common form. The upper, superficial; the lower, deep.

VIII. Agar plate, eight days at 37° . $\times 50$. Larger colony superficial; smaller colonies, deep.

IX. Microscopic preparation from a two days' bouillon culture at 37° . \times 700. The individual cocci are usually more regularly round.

X. Microscopic preparation from a two days' agar culture. Shorter chains. \times 1000.

XI. Microscopic preparation, designated Streptococcus conglomeratus. Smear preparation from the blood of the spleen from a case of scarlatina. Copied from Kurth (Kaiserl. Gesundheitsamt., Bd. VII, cfr. xy, 6 and 8).

XII. Chains of streptococci, before and during division. Highly magnified.



PLATE 2.

Streptococcus lanceolatus. Gamaleia. (Diplococcus pneumonia: A. Fränkel.) (Pneumococcus.)

I. Gelatin stab culture, ten days at 22°.

II. Agar streak culture, four days at 37°.

III. Agar stab culture, four days at 37°. Stab canal.

IV. Agar stab culture, four days at 37°. Surface growth.

V. Agar plate, three days at 37°. Natural size.

VI. Agar plate, three days at 37°. \times 50. Superficial colony. The darker colony lies near the surface.

VII. Agar plate, three days at 37°. \times 50. Deep colonies.

VIII. Gelatin plate, eight days at 22°. The upper colony is superficial; the lower ones, deep.

IX. Smear preparation from pneumonic sputum. \times 1000.

X. Pure culture from a three days' old agar plate. \times 1000.

XI. Microscopic preparations: (a) Diplococci as single pairs and in chains. Highly magnified. (b) Diplococci surrounded by gelatinous capsules.



XÍ.